

RESULTS OF THE CLASSIFICATION OF SUPERCONFORMAL ALGEBRAS IN TWO DIMENSIONS

E.S. Fradkin* and **V.Ya. Linetsky**

Department of Theoretical Physics,
P.N.Lebedev Physical Institute, Leninsky pr. 53, Moscow 117924

Abstract

A list of superconformal chiral operator product expansion algebras with quadratic nonlinearity in two dimensions is completed on the basis of the known classification of little conformal Lie superalgebras. In addition to the previously known cases and the constructed in our previous paper exceptional $N = 8$ superalgebra associated with $F(4)$, a novel exceptional $N = 7$ superconformal algebra associated with $G(3)$ is found, as well as a whole family of superalgebras containing affine $\widehat{su}_2 \oplus \widehat{usp}_{2N}$. A classification scheme for quasisuperconformal algebras is also outlined.

February 1992

* present address: Institut für Theoretische Physik, ETH-Hönggerberg, CH-8093 Zürich

Recently we have found a novel $N = 8$ superconformal algebra in two dimensions [1] with supercurrents in the eight-dimensional spinor representation of $\text{Spin}(7)$. It involves a peculiar composite $:JJ:$ term in the anticommutator of two supercurrents, similar to the superconformal algebras of Refs.[2-5]. It has encouraged us to see whether there exist any other previously unknown superconformal algebras involving quadratic composites (simple superconformal algebras of the Lie type involving operators with scale dimensions 2, $3/2$, 1 and $1/2$ have been classified by Ramond and Schwarz [6]). As a result we have completed a list of superconformal algebras of this category which is given in the present letter. Here we present only the final results, all the proofs and details being postponed to a more detailed paper [7].

Suppose we are given a pair (g, ρ) of a compact Lie algebra g and its representation ρ . Consider a two-dimensional conformal field theory with an energy-momentum $T(z)$, a multiplet of supercurrents $G^\alpha(z)$ in the representation ρ of g and a multiplet of Kac-Moody currents $J^A(z)$ in the adjoint representation of g , scale dimensions being 2, $3/2$ and 1, respectively. (We are concerned with the holomorphic sector only.)

Then a problem we address is to enumerate all the admissible pairs (g, ρ) such that the corresponding associative $\mathbf{Z}_2$ -graded chiral operator product expansion closes on T , G , J and their (normal ordered) composites only, and to evaluate explicitly the corresponding superconformal algebras. We assume the resulted superconformal algebras, which we sometimes denote $SC(g, \rho)$, to be simple in the following sense. Consider an associative algebra U with generating elements $\{L_n, G_r^\alpha, J_n^A\}$ (the modes of $T(z)$, $G^\alpha(z)$ and $J^A(z)$) and defining relations determined by the corresponding OPE. Then a SC is said to be simple iff U contains no non-trivial ideals generated by a non-trivial set of elements linear in generating elements. In case when SC is of the Lie-type (*i.e.* involvs no composites in the supercommutators) this definition is equivalent to the usual definition of simplicity of a Lie superalgebra and U is its universal enveloping.

Actually, we have proved that there exists a simple $SC(g, \rho)$ iff there exists a simple little (finite-dimensional) conformal Lie superalgebra $LSC(g, \rho)$ with its even subalgebra isomorphic to $\mathfrak{sl}(2; \mathbf{R}) \oplus g$ and its odd subspace transforming in the representation $(2, \rho)$ of $\mathfrak{sl}(2; \mathbf{R}) \oplus g$. This proposition is quite non-trivial since the LSC is not generally, *i.e.* when composites are presented, a subalgebra of SC (the situation here is similar to, *e.g.*, the relation of $\mathfrak{sl}_N$ to W_N) and requires a proof based on the detailed analysis of the Jacobi identities for the quantum algebra SC . (For how to extract LSC from SC in non-linear case see Ref.[1].) However, we will not discuss it here but rather directly pass to the list of $SC(g, \rho)$ resulted from the above proposition. It is based on the list of little conformal Lie superalgebras given in Refs.[8], [9] (which in turn is based on Kac's classification of simple Lie superalgebras). There are three classical series, one

continuous family and two exceptional cases related with octonions.

- Case *I* : $osp(N|2; \mathbf{R})$, $N = 1, 2, \dots$, $g = so(N)$, $\rho = N$.

For $N = 1$ the corresponding OPE defines the Neveu-Schwarz-Ramond superalgebra [10], [11], for $N = 2$ it is the superalgebra of Ademollo et al. [12] involving an $so(2)$, and for $N \geq 3$ $SC(so(N), N)$ has been constructed by Knizhnik [2] and by Bershadsky [3]. Starting at $N = 3$ in order to close the OPE one has to introduce the quadratic non-linearity $:JJ:$ in the right-hand side of $G^i(z)G^j(w)$. In cases $N = 3$ and $N = 4$ it is shown by Goddard and Schwimmer [4] that the corresponding non-Lie type OPEs can be obtained by factoring out free fermions and bosons with scale dimensions $1/2$ and 0 from the $N = 3$ and $N = 4$ Lie superalgebras of Refs.[12].

- Case *II* : $su(1, 1|N)$, $N = 3, 4, \dots$, $g = su(N) \oplus u(1)$, $\rho = N \oplus \bar{N}$, and $su(1, 1|2)/u(1)$, $g = su(2)$, $\rho = 2 \oplus \bar{2}$.

While in the later case the superconformal algebra involves an $\widehat{su}(2)$ Kac-Moody algebra and closes as a Lie superalgebra [12], for all $N \geq 3$ one has to introduce quadratic composites [2], [3]. Then the Jacobi identities require the central charges for the affine $\widehat{su}(N)$ and $\hat{u}(1)$ to be of opposite signs that generally leads to non-unitarity of the Verma modules.

- Case *III* : $osp(4^*|2M)$, $M = 1, 2, \dots$, $g = usp(2M) \oplus su(2)$, $\rho = (2M, 2)$.

Meanwhile the case $M = 1$ is isomorphic to the subcase $N = 4$ of the case I, a whole family of superconformal algebras SC with $M = 2, 3, \dots$ seems to be not constructed previously. So we present our results for this case.

Let $\alpha, \beta, \gamma, \delta = 1, 2$ and $A, B, C, D = 1, 2, \dots, 2M$ ($M \geq 2$) be indices in the fundamental representation of $usp(2)$ ($\simeq su(2)$) and $usp(2M)$, respectively. They are raised and lowered with the help of the symplectic metrics $\varepsilon^{\alpha\beta} = -\varepsilon^{\beta\alpha}$, $\varepsilon_{\alpha\beta} = -\varepsilon_{\beta\alpha}$, $\varepsilon_{\alpha\gamma}\varepsilon^{\beta\gamma} = \delta_{\alpha}^{\beta}$ and $E^{AB} = -E^{BA}$, $E_{AB} = -E_{BA}$, $E_{AC}E^{BC} = \delta_A^B$, respectively. Then a SC is given by the following supercommutation relations among the Virasoro L_n (associated to $sl(2; \mathbf{R})$ subalgebra of $osp(4^*|2M)$), $\widehat{su}(2) \oplus \widehat{usp}(2M)$ Kac-Moody currents $J_m^{\alpha\beta} = J_m^{\beta\alpha}$ and $J_m^{AB} = J_m^{BA}$ (associated to g) and supercurrents $G_r^{\alpha A}$ (associated to ρ)

$$[L_m, L_n] = (m - n)L_{m+n} + \frac{c}{12}m(m^2 - 1)\delta_{m+n,0}, \quad (1a)$$

$$\begin{aligned} [J_m^{\alpha\beta}, J_n^{\gamma\delta}] &= \varepsilon^{\beta\gamma}J_{m+n}^{\alpha\delta} + \varepsilon^{\alpha\delta}J_{m+n}^{\beta\gamma} + \varepsilon^{\alpha\gamma}J_{m+n}^{\beta\delta} + \varepsilon^{\beta\delta}J_{m+n}^{\alpha\gamma} \\ &\quad - k_2(\varepsilon^{\alpha\gamma}\varepsilon^{\beta\delta} + \varepsilon^{\beta\gamma}\varepsilon^{\alpha\delta})m\delta_{m+n,0}, \end{aligned} \quad (1b)$$

$$[J_m^{AB}, J_n^{CD}] = E^{BC}J_{m+n}^{AD} + E^{AD}J_{m+n}^{BC} + E^{AC}J_{m+n}^{DB} + E^{BD}J_{m+n}^{CA}$$

$$-k_1(E^{AC}E^{BD} + E^{BC}E^{AD})m\delta_{m+n,0}, \quad (1c)$$

$$[L_m, J_n^{\alpha\beta}] = -nJ_{m+n}^{\alpha\beta}, \quad [L_m, J_n^{AB}] = -nJ_{m+n}^{AB}, \quad (1d)$$

$$[L_m, G_r^{\alpha A}] = \left(\frac{m}{2} - r\right)G_{m+r}^{\alpha A}, \quad (1e)$$

$$[J_m^{\alpha\beta}, G_r^{\gamma A}] = \varepsilon^{\beta\gamma}G_{m+r}^{\alpha A} + \varepsilon^{\alpha\gamma}G_{m+r}^{\beta A} \quad (1f),$$

$$[J_m^{AB}, G_r^{\alpha C}] = E^{BC}G_{m+r}^{\alpha A} + E^{AC}G_{m+r}^{\alpha B}, \quad (1g)$$

$$\begin{aligned} \{G_r^{\alpha A}, G_s^{\beta B}\} &= 2\varepsilon^{\alpha\beta}E^{AB}L_{r+s} + (r-s)[\sigma_1\varepsilon^{\alpha\beta}J_{r+s}^{AB} + \sigma_2E^{AB}J_{r+s}^{\alpha\beta}] \\ &\quad + b\left(r^2 - \frac{1}{4}\right)\varepsilon^{\alpha\beta}E^{AB}\delta_{r+s,0} + \gamma(JJ)_{r+s}^{\alpha A, \beta B}, \end{aligned} \quad (1h)$$

where the normal ordered quadratic term reads as follows

$$\begin{aligned} (JJ)_m^{\alpha A, \beta B} &= 2\varepsilon^{\alpha\beta}E^{AB}\{\varepsilon_{\alpha_1\beta_1}\varepsilon_{\alpha_2\beta_2}(J^{\alpha_1\alpha_2}J^{\beta_1\beta_2})_m - E_{A_1B_1}E_{A_2B_2}(J^{A_1A_2}J^{B_1B_2})_m\} \\ &\quad + 8(J^{\alpha\beta}J^{AB})_m + 4\varepsilon^{\alpha\beta}E_{CD}\{(J^{AC}J^{BD})_m - (J^{BC}J^{AD})_m\}, \end{aligned} \quad (2a)$$

where

$$(J^{AB}J^{CD})_m = \sum_{n \in \mathbf{Z}} : J_n^{AB}J_{m-n}^{CD} : \quad (2b)$$

and similar for $(J^{\alpha\beta}J^{\gamma\delta})_m$. The central charges k_1 , k_2 , c and b and the structure constants σ_1 , σ_2 and γ are fixed by the Jacobi identities to be

$$k_1 = -(k + 2M + 4)/2, k_2 = k, \quad (3a)$$

$$b = -\frac{k(k + 2M + 4)}{k - 2M + 4}, \quad (3b)$$

$$c = -\frac{3k(k + 2M + 4)}{k - 2M + 4} + \frac{6k + (2M + 1)(M - 2)(k + 2M + 4)}{k - 2M + 4}, \quad (3c)$$

$$\sigma_1 = -\frac{2k}{k-2M+4}, \quad \sigma_2 = \frac{k+2M+4}{k-2M+4}, \quad (3d)$$

$$\gamma = \frac{1}{2(k-2M+4)}. \quad (3e)$$

Here k is the only free parameter which determines the values of all the central charges. One sees k_1 and k_2 have opposite signs that leads to non-unitarity similar to the case II (for $N \geq 3$). To check up on the Jacobi a crucial part is played by the Fierz-type identity

$$\varepsilon^{\alpha\beta}\delta_\delta^\gamma + \varepsilon^{\beta\gamma}\delta_\delta^\alpha + \varepsilon^{\gamma\alpha}\delta_\delta^\beta = 0 \quad (4)$$

which follows from the fact that the antisymmetrization in three two-valued indices gives zero.

- Case IV: $D^1(2, 1; \alpha)$, $g = su(2) \oplus su(2)$, $\rho = (2, 2)$ (one-parameter family; $\alpha \rightarrow 1/\alpha$ gives an isomorphism).

For $M = 1$ the same identity (4) holds also for capital latin indices A, B, C, D and there appears a possibility to have got one more free parameter, say α , in the algebra (1). Then two Kac-Moody central charges k_1 and k_2 for "left" and "right" affine $\widehat{su}(2)$ become independent (in fact α governs a "left-right asymmetry") and it makes it possible to have both k_1 and k_2 simultaneously positive integer. The resulting non-linear superconformal algebra is obtained in Ref.[4] by factoring out free fermions and bosons from the Lie superalgebra of Refs.[13]. The unitary highest weight representations are studied and applied to superstring compactification in Ref.[5].

- Case V : $G(3)$, $g = G_2$, $\rho = 7$.

Consider an octonionic algebra $\mathbf{O}$ with seven imaginary units e^a , $a = 1, 2, \dots, 7$, and a composition law

$$e^a e^b = -\delta^{ab} + \Gamma^{abc} e^c, \quad (5)$$

where Γ^{abc} is a totally skew G_2 -invariant tensor. Thanks to the alternativity of $\mathbf{O}$, Γ obeys a crucial identity

$$\Gamma^{abe}\Gamma^{cde} + \Gamma^{cbe}\Gamma^{ade} = 2\delta^{ac}\delta^{bd} - \delta^{ad}\delta^{bc} - \delta^{ab}\delta^{cd}. \quad (6)$$

An exceptional $N = 7$ superconformal algebra $SC(G_2, 7)$ involves the Virasoro L_m , $\hat{G}_2$ Kac-Moody currents $J_m^{ab} = -J_m^{ba}$ satisfying the property (actually it defines the projection onto $G_2 \subset so(7)$; for the properties of G_2 see *e.g.* [14], or [15]-[17] where the supergroups $G(3)$ and $F(4)$ are described in detail)

$$\Gamma^{abc} J_m^{bc} = 0, \quad (7)$$

and supercurrents G_r^a in the fundamental seven-dimensional representation of G_2 . Then our result for the superalgebra reads as follows (we omit the obvious commutation relations with L_m)

$$[J_m^{ab}, J_n^{cd}] = \delta^{bc} J_{m+n}^{ad} + \delta^{ad} J_{m+n}^{bc} - \delta^{ac} J_{m+n}^{bd} - \delta^{bd} J_{m+n}^{ac} + \frac{1}{3} \Gamma^{abe} \Gamma^{cdf} J_{m+n}^{ef} - k(\delta^{ac} \delta^{bd} - \delta^{ad} \delta^{bc} - \frac{1}{3} \Gamma^{abe} \Gamma^{cde}) m \delta_{m+n,0} \quad (8a)$$

$$[J_m^{ab}, G_r^c] = \delta^{bc} G_{m+r}^a - \delta^{ac} G_{m+r}^b + \frac{1}{3} \Gamma^{abe} \Gamma^{cde} G_{m+r}^d, \quad (8b)$$

$$\{G_r^a, G_s^b\} = 2\delta^{ab} L_{r+s} + \sigma(r-s) J_{r+s}^{ab} + b(r^2 - \frac{1}{4}) \delta^{ab} \delta_{r+s,0} + \gamma P_{cd,ef}^{ab} (J^{cd} J^{ef})_{r+s}, \quad (8c)$$

where the tensor P is uniquely determined from the Jacobi identities

$$P_{ab,cd} = \{\lambda_{ab}, \lambda_{cd}\} + \frac{4}{3}(\delta_{ac} \delta_{bd} - \delta_{ad} \delta_{bc} - \frac{1}{3} \Gamma_{abe} \Gamma_{cde}) \mathbf{1}, \quad (9)$$

and λ_{ab} are G_2 matrices in the fundamental representation

$$\lambda_{ab,cd} = \delta_{ac} \delta_{bd} - \delta_{ad} \delta_{bc} - \frac{1}{3} \Gamma_{abe} \Gamma_{cde}. \quad (10)$$

All the constant parameters c, b, σ and γ are uniquely fixed by associativity to be the following rational functions of the $\hat{G}_2$ Kac-Moody level k :

$$c = \frac{9k}{2} + \frac{2k}{k+3}, \quad (11a)$$

$$b = k\sigma = \frac{3k}{2} (1 - \frac{4}{3(k+3)}), \quad (11b)$$

$$\sigma = \frac{3}{2} (1 - \frac{4}{3(k+3)}), \quad (11c)$$

$$\gamma = \frac{3}{16(k+3)}. \quad (11d)$$

This time a crucial role in checking up the Jacobi is played by the octonionic identity (6).

There exists a natural embedding of the $N = 1(NSR)$ superconformal algebra in our exceptional $N = 7$ algebra (more detailed discussion for the $N = 8$ exceptional case see in Ref.[1]). Choose the supergenerator G_r of the $N = 1$ superalgebra we are about to construct to be equal to G_r^1 and redefine the Virasoro generators according to

$$\tilde{L}_m = L_m + \frac{\gamma}{2} P_{ab,cd}^{11} (J^{ab} J^{cd})_m. \quad (12)$$

Then it is easy to see that $\tilde{L}_m$ and G_r form an $N = 1$ superalgebra with the Virasoro central charge

$$\tilde{c} = \frac{9k}{2} \left(1 - \frac{4}{3(k+3)}\right). \quad (13)$$

The structure of new Virasoro generators (12) is reminiscent of the GKO coset construction for $G_2/su(3)$.

Note that central charges $c, \tilde{c}$ and b are positive when the Kac-Moody level k is positive integer, the obvious necessary condition for unitarity thus being satisfied. The same also holds for the next case VI.

- Case VI : $F(4)$, $g = Spin(7)$, $\rho = 8_s$.

The exceptional $N = 8$ conformal superalgebra $SC(Spin(7), 8_s)$ associated with $F(4)$ has been constructed in our previous paper [1].

To summarize, in our classification of superconformal OPEs in two dimensions there are three classical families, one continuous family and two exceptional cases with $N = 7$ and $N = 8$. Of special interest are the problems of constructing representations and field-theoretic realizations of the exceptional algebras. Probably they could be obtained via the quantum Drinfeld-Sokolov reduction based on the underlying little superalgebras $G(3)$ and $F(4)$.

Note Added

In this paper we have considered superconformal algebras with the standard spin-statistics relation. Recently so-called quasisuperconformal algebras have appeared in the context of the quantum Drinfeld-Sokolov reduction and related approaches [16]-[18]. They involve bosonic currents of dimension $3/2$ in a representation ρ of some Lie algebra g along with the energy-momentum and $\hat{g}$ KM currents. A list of possible pairs (g, ρ) may be completed on the basis of Cartan's classification of symmetric subalgebras of simple Lie algebras. One requires the symmetric subalgebra to be isomorphic to $su(2) \oplus g$ and the supplementary subspace to transform in its representation $(2, \rho)$ with $\rho^{\mathbf{C}}$ possessing a $g^{\mathbf{C}}$ -invariant symplectic metric. As a result one finds eight relevant cases (g, ρ and underlying simple Lie algebras are presented): Case I: $u(N), N \oplus \bar{N}, su(N+2)$; Case II: $sp(2M), 2M, sp(2M+2)$; Case III: $su(2) \oplus so(N), (2, N), so(N+4)$; Case IV: $su(2), \dim \rho = 4, G_2$; Case V : $sp(6), \dim \rho = 14, F_4$; Case VI: $su(6), \rho = 20, E_6$; Case VII: $so(12), \rho = 32', E_7$; Case VIII : $E_7, \rho = 56, E_8$. Quasisuperconformal algebras corresponding to the Cases I and II have been studied in Refs.[16] and [17]. The Case III represents a bosonic counterpart of our Case III superconformal algebras based on $osp(4^*|2M)$. Five exceptional cases will be considered in a separate publication. However it should be stressed that there are difficulties in the Cases *II–VIII* with imposing the hermiticity condition, since representations ρ are pseudoreal and, in particular, it is not possible to define real Euclidean $su(2)$ spinors.

Then two series of $\mathbf{Z}_2 \times \mathbf{Z}_2$ -graded superconformal algebras have been introduced in Ref.[18]. They contain affine superalgebras $\widehat{su}(m|n)$ and $\widehat{osp}(N|2M)$ and in fact combine superconformal and quasisuperconformal algebras together. Our results suggest the existence of a third series of this type combining the Cases III from our two lists. It contains affine $\widehat{su}(2) \oplus \widehat{osp}(N|2M)$.

Acknowledgement We are grateful to W.Troost for sending us a copy of the preprint [18]. E.S. Fradkin thanks Prof. J. Fröhlich and the “Zentrum für Theoretische Studien” at ETH for hospitality and also the SNF (Swiss National Fonds) for financial support.

References

1. E.S.Fradkin and V.Ya.Linetsky, *ICTP/91/348* (Phys. Lett. *B*, to be published).
2. V.G.Knizhnik, Theor. Math. Phys. 66 (1986) 68.
3. M.Bershadsky, Phys. Lett. B174 (1986) 285.
4. P.Goddard and A.Schwimmer, Phys.Lett. B214 (1988) 209.
5. *M.Günaydin*, J.L.Petersen, A.Taormina and A.Van Proeyen, Nucl. Phys. B322 (1989) 402.
6. P.Ramond and J.H.Schwarz, Phys. Lett. B65 (1976) 75.
7. E.S.Fradkin and V.Ya.Linetsky, Classification of superconformal algebras in two dimensions, in preparation.
8. W.Nahm, Nucl. Phys. B135 (1978) 149.
9. *M.Günaydin*, G.Sierra and P.K.Townsend, Nucl. Phys. B274 (1986) 429.
10. P.Ramond, Phys. Rev. D3 (1971) 2415.
11. A.Neveu and J.H.Schwarz, Nucl. Phys. B31 (1971) 86.
12. M.Ademollo et al., Phys. Lett. B62 (1976) 105; Nucl. Phys. B111 (1976) 77; B114 (1976) 297.
13. A.Sevrin, W.Troost and A.Van Proeyen, Phys. Lett. B208 (1988) 447; K.Schoutens, Nucl. Phys. B295 (1988) 634.
14. R.E.Behrends, J.Dreitlein, C.Fronsdal and W.Lee, Rev. Mod. Phys. 34 (1962) 5.
15. M.Scheunert, W.Nahm and V.Rittenberg, J. Math. Phys. 17 (1976) 1640; B.S. De Witt and P. van Nieuwenhuizen, J. Math. Phys. 23 (1982) 1953; A.Sudbery, J. Math. Phys. 24 (1983) 1986.
16. A.M.Polyakov, in "Physics and Mathematics of Strings", World Scientific, 1990; M.Bershadsky, Commun. Math. Phys. 139 (1991) 71; F.A.Bais, T.Tjin and P. van Driel, Nucl. Phys. B357 (1991) 632; A.Bilal, CERN-TH.6294/91.
17. L.J.Romans, Nucl. Phys. B357 (1991) 549; F.Defever, Ph.D.Thesis, Leuven 1991.
18. F.Defever, W.Troost and Z.Hasiewicz, Preprint KUL-TF-91-26&ITP-UWr91-779 (Phys. Lett. B273 (1991) 51).